

Food and Drug Administration, HHS

§ 558.95

§ 558.78 Bacitracin zinc.

(a) *Specifications.* Type A medicated articles containing bacitracin zinc equivalent to 10, 25, 40, or 50 grams per pound bacitracin.

(b) *Approvals.* See No. 054771 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.70 of this chapter.

(d) *Conditions of use.* (1) It is used as follows:

Bacitracin zinc in grams per ton	Combinations in grams per ton	Indications for use	Limitations	Sponsor
(i) 4 to 50	Chickens: for increased rate of weight gain and improved feed efficiency.	Growing chickens	054771
(ii) 4 to 50	Turkeys and pheasants: for increased rate of weight gain and improved feed efficiency.	Growing turkeys and pheasants	054771
(iii) 5 to 20	Quail; for increased rate of weight gain and improved feed efficiency.	Growing quail; feed as the Type C feed to starting quail through 5 weeks of age.	054771
(iv) 10 to 25	Laying chickens; improved feed efficiency and increased egg production.	054771
(v) 10 to 50	Swine; increased rate of weight gain and improved feed efficiency.	Growing and finishing swine	054771
(vi) 20	Growing-finishing swine; increased rate of weight gain.	In Type C feed	054771
(vii) 20 to 40	Growing-finishing swine; improved feed efficiency.do	054771

(2) It is used in feed for growing cattle at 35 to 70 milligrams per head per day as follows:

(i) To aid in stimulating growth and improving feed efficiency.

(ii) For increased rate of weight gain and improved feed efficiency; see sponsor 054771.

(3) Bacitracin zinc may also be used in combination with:

(i) Amprolium and ethopabate as in § 558.58.

(ii) Clopidol as in § 558.175.

(iii) Decoquinat as in § 558.195.

(iv) Lasalocid as in § 558.311.

(v) Monensin as in § 558.355.

(vi) Naracin as in § 558.363.

(vii) [Reserved]

(viii) Robenidine as in § 558.515.

(ix) Salinomycin as in § 558.550.

[41 FR 10994, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.78, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and at www.fdsys.gov.

§ 558.95 Bambermycins.

(a) *Approvals.* See sponsors in § 510.600(c) of this chapter for use of Type A medicated articles as in paragraph (d) of this section:

(1) No. 016592: 2, 4, and 10 grams per pound for use as in paragraphs (d)(1), (d)(2), (d)(3), and (d)(4) of this section.

(2) No. 012286: 2 grams for use as in paragraph (d)(2) of this section and 0.4 and 2 grams per pound for use as in paragraph (d)(3).

(b) *Special considerations.* (1) Bambermycins liquid Type B feeds may be manufactured from dry bambermycins Type A articles. The liquid Type B feeds must have a pH of 3.8 to 7.5, moisture content of 30 to 45 percent.

(2) The expiration date for the liquid Type B feed is 8 weeks after date of manufacture. The expiration date for the dry Type C feed made from the liquid Type B feed is 1 week after date of manufacture.

(c) [Reserved]

(d) *Conditions of use*—(1) *Chickens.* Use in medicated feed as follows: